



Client		FARMS	Produc	t: CGF-2020)-01	Date F	Reported:	08/20/20	
	Attn: BILL LONGBRAKE			Date Sample	ed: 08/04/20	Labor	atory #	C20-770	
	1201 S Grover Lane			Date Receive	ed: 08/06/20	Rev	eiwed by Br	rent Thyssen,	CPSSc
	Palmer, AK 99645				Invoice #:	C20-770	PO#:		
	907-863-0802					A	mount:	\$300.00	
				Nutrien	ts				
	Method	As Received	Dry Wt.	Units	Low	Normal	Hi	igh	Typical Range
Moisture	70 C	35		%	*****				15 to 40
Solids	70 C	65		%	*****	****			60 to 85
рН	1:5	7.4	NA	SU	*****	****			5.5 to 8.5
E.C. (Sol. Salts)	1:5	3.08	4.75	mmhos/cm	*****	****			below 5.0
Total N	TMECC 04.02D	0.91	1.41	%	*****	*			1 to 5
Organic C	TMECC 04.01A	10.2	15.8	%	*****				18 to 45
Organic Matter	TMECC 05.07A	19.5	30.1	%	*****				40 to 60
Ash	550 C	45.2	69.9	%	*****	****	**		40 to 60
Ammonium -N	TMECC 05.02C	25	39	mg/kg	****				90 to 450
Nitrate-N	TMECC 04.02B	272	420	mg/kg	*****	****	k		50 to 250
Chloride	TMECC 04.12D	1704	2632	mg/kg	*****	****			500 to 5000
Sulfate-S	TMECC 04.12D	91	140	mg/kg					
CaCO ₃	TMECC 04.08A	19	30	lbs/T	*****				20 to 80
Phosphorous	TMECC 04.12B/04.14A	0.25	0.38	%					
P ₂ O ₅	calculation	0.57	0.88	%	****				1 to 8
Potassium	TMECC 04.12B/04.14A	0.88	1.36	%					
K₂O	calculation	1.06	1.64	%	****				3 to 12
Calcium	TMECC 04.12B/04.14A	0.73	1.1	%	*****	*			0.5 to 10
Magnesium	TMECC 04.12B/04.14A	0.47	0.73	%	*****	*****			0.05 to 0.7
Sodium	TMECC 04.12B/04.14A	0.05	0.08	%	*****				0.05 to 0.7
Sulfur	TMECC 04.12B/04.14A	0.13	0.20	%	*****	****			0.1 to 1.0
Boron	TMECC 04.12B/04.14A	1	2	mg/kg	***				25 to 150
Zinc	TMECC 04.12B/04.14A	69	106	mg/kg	*****	*			100 to 600
Manganese	TMECC 04.12B/04.14A	346	535	mg/kg	*****	****			250 to 750
Copper	TMECC 04.12B/04.14A	21	32	mg/kg	***				100 to 500
Iron	TMECC 04.12B/04.14A	13753	21240	mg/kg	*****	****			1000 to 25000
C/N ratio			11	ratio	*****				18 to 24
C/P Ratio			41	ratio	*****				80 to 140
Ag Index			9	ratio	*****	****			3 to 10

Respiration & Stability

			nespiration o	Clubinty			
	Method		Units	Low	Normal	High	Normal
CO2 Evolution	TMECC 05.08	0.2	mg CO ₂ -C/g OM/day	**			1 to 7
_	TMECC 05.08	0.2	mg CO ₂ -C/g TS/day	****			0.5 to 5
Stability Rating		Very Stable					

Sample was received, handled and tested in accordance with TMECC procedures

Page 1



COUNTRY GARDEN FARMS	Product: CGF-2020-01	Date Reported: 08/20/20
1201 S Grover Lane	Date Sampled: 08/04/20	Laboratory # C20-770
Palmer, AK 99645	Date Received: 08/06/20	Reveiwed by Brent Thyssen, CPSSc
907-863-0802		
	1201 S Grover Lane Palmer, AK 99645	1201 S Grover LaneDate Sampled:08/04/20Palmer, AK 99645Date Received:08/06/20

	Cucumber Bioassay									
	Method		Units	Low	Normal	Normal				
Emergence	TMECC 05.05A	100	%	*******************		80 to 100				
Vigor	TMECC 05.05A	93	%	**********		85 to 100				
Maturity	V	ery Mature: safe for use i	n containe	rs						

Pathogens										
			Date Tested	8/11/2020				_		
	Method		units		Low	No	rmal		High	Normal
Fecal Coliforms	TMECC 07.01AB	Not Tested	MPN/g							Less than 1000
Salmonella	TMECC 07.02A	ND	MPN/4g	Pass	*					Less than 3
		ND = None Detected	d Fe	ecal Coliforms MI	DL 3.6	MPN/g	Salmor	nella MDL	1	MPN/4g

EPA 503 Metals								
	Method	Dry Wt.	Units	Low	Normal	High	MDL	EPA Limit
Arsenic	TMECC 04.12B/04.14A	9.3	mg/kg	******			0.30	41
Cadmium	TMECC 04.12B/04.14A	<mdl< td=""><td>mg/kg</td><td></td><td></td><td></td><td>0.08</td><td>39</td></mdl<>	mg/kg				0.08	39
Chromium	TMECC 04.12B/04.14A	21.5	mg/kg				0.09	-
Cobalt	TMECC 04.12B/04.14A	8.0	mg/kg	****			0.09	1200
Copper	TMECC 04.12B/04.14A	31.7	mg/kg	****			0.25	1500
Mercury	TMECC 04.12B/04.14A	0.04	mg/kg	****			0.002	17
Molybdenum	TMECC 04.12B/04.14A	3.7	mg/kg	****			0.17	75
Nickel	TMECC 04.12B/04.14A	17.2	mg/kg	****			0.12	420
Lead	TMECC 04.12B/04.14A	2.5	mg/kg	****			0.23	300
Selenium	TMECC 04.12B/04.14A	<mdl< td=""><td>mg/kg</td><td></td><td></td><td></td><td>0.97</td><td>100</td></mdl<>	mg/kg				0.97	100
Zinc	TMECC 04.12B/04.14A	106	mg/kg	****			0.25	2800
	Metals Assay	Pass						

Particle Size Distribution TMECC 2.02 B & C

3 76.2 100 2 50 100 Total Plastic 0.00 1 25 100 Film Plastic 0.00 3/4 19.1 100 Glass 0.00 5/8 16 100 Metal 0.00 1/2 12.5 100 Sharps 0.00 3/8 9.5 98 Contract Contract	inches	mm	% Passing	Inerts	% by wt.
1 25 100 Film Plastic 0.00 3/4 19.1 100 Glass 0.00 5/8 16 100 Metal 0.00 1/2 12.5 100 Sharps 0.00 3/8 9.5 98 100 100	3	76.2	100		
3/4 19.1 100 Glass 0.00 5/8 16 100 Metal 0.00 1/2 12.5 100 Sharps 0.00 3/8 9.5 98 100 100	2	50	100	Total Plastic	0.00
5/8 16 100 Metal 0.00 1/2 12.5 100 Sharps 0.00 3/8 9.5 98	1	25	100	Film Plastic	0.00
1/2 12.5 100 Sharps 0.00 3/8 9.5 98	3/4	19.1	100	Glass	0.00
3/8 9.5 98	5/8	16	100	Metal	0.00
	1/2	12.5	100	Sharps	0.00
	3/8	9.5	98		
1/4 6.3 91	1/4	6.3	91		

Sample was received, handled and tested in accordance with TMECC procedures

Page 2



COUNTRY GARDEN FARMS Attn: BILL LONGBRAKE 1201 S Grover Lane Palmer, AK 99645 907-863-0802

DATE REC 6-Aug-20 INVOICE # 6-Aug-20 LAB # C20-770 Date Reported: 08/20/20

NUTRIENT REPORT

SAMPLE I.D.:	CGF-2020-01				
As Received:	<u>%SOLIDS</u> 64.75		<u>%WATER</u> 35.25		
TOTAL	100%D			AS RECE	
ELEMENTS	100%D	lbs/ton		A3 NECE %	lbs/ton
TN	1.41	28.20		0.91	18.3
Р	0.38	7.68		0.25	5.0
P205	0.88	17.65		0.57	11.4
К	1.36	27.28		0.88	17.7
K20	1.64	32.74		1.06	21.2
S	0.20	3.92		0.13	2.5
Ca	1.13	22.5		0.73	14.6
Mg	0.73	14.55		0.47	9.4
Na	0.08	1.60		0.05	1.0
С	15.80	316		10.2	205
	mg/kg	lbs/ton		mg/kg	lbs/ton
Zn	106	0.21		69	0.14
Mn	535	1.07		346	0.69
Cu	32	0.06		21	0.04
Fe	21240	42.48		13753	27.5
В	2	0.00		1.10	0.00
Nitrate N	420	0.84		272.0	0.54
Ammonium N	39	0.08		25	0.05
C:N Ratio				11	
pH				7.4	
E.C.	4.75			3.08	



Client: COUNTRY GARDEN FARMS Product: CGF-2020-01 C20-770 l ah #

INTERPRETATION GUIDE

Date Sampled: 08/04/20 08/06/20 Date Received: Date Reported: 08/20/20

SAFETY INTERPRETATIONS

Pathogens

Fecal coliform bacteria are present in the gut and fecal mater of warm-blooded animals. Their presence is used as an indicator of the presence of possible human pathogens. The heat generated during proper composting is lethal to fecal coliform and other human pathogens. A test value below 1,000 per gram of compost is considered generally safe for human contact. As the compost is stored or transported, the temperature is no longer lethal for coliform bacteria and there is the possibility for regrowth or contamination by birds or other animals. Your compost was not tested for fecal coliform.

Salmonella is a human pathogenic bacteria and a good indicator of other human pathogens. It is regularly used to monitor the liklihood of human pathogen presence in biosolids VERY SAFE

Your compost was tested for salmonella bacteria and found to be:

Heavy Metals

9 heavy metals were identified with maximum concentration limits for land application in biosolids by USEPA in 40 CFR Part 503,B. Ongoing applications to the land are prohibited if any metal concentration exceed the limits in Table 3 of Part 503.13. If the bars on the "Heavy Metals" for your compost are within or below the "Normal" range, your compost is safe

to use as a soil amendment.

COMPOST STABILITY AND MATURITY

Respiration

Respiration is the measurement of microbially generated CO2 from the compost when incubated at optimal temperature and moisture. It provides an indication of whether the composting process is complete and whether the compost is mature and ready for use. However, other factors may be limiting microbial activity (see C:N Ratio below)

Your Compost was rated as Very Stable: well cured, finished compost; no odors or plant toxicity

Maturity

Bioassav

Cucumbers are grown in a fixed blend of your compost and a commercial potting mix maintained at optimum moisture and temperature. Cucumbers are relatively insensitive to salinity, but very sensative to ammonia, organic acids and herbicide residue. Emergence and Vigor are rated: results greater than 80% indicate that your compost is mature and/or contains no hervicide carryover. Very high salinity can also reduce assay results.

100

Your Compost Emergence %

Your Compost vigor %

93

Total Nitrogen, Nitrate & Ammonium

Ammonia is produced as a gas in the early stages of composting. The ammonium is nitrified to nitrate as the compost matures. Ammonia is toxic to plants at relatively low concentrations but under moist conditions is converted to ammonium which is less toxic. Nitrate is not toxic, but does contribute to overall salinity if very high. The pH of the compost typically starts out low as organic acids are released, then increases as ammonia is produced, then settles back towards nuetral (7.0) as ammonium is nitrified and the compost matures.

Your Compost Ammonium level was	39	Your Compost Ammonium:Nitrate ratio was	0
Your Compost Ammonium: Total N ratio was	0.00	Your Compost pH was	7.4

Considering all the factors above, your Compost is Very Mature: safe for use in containers

FERTILITY INTERPRETATIONS

C:N Ratio

The carbon to nitrogen ratio is important to determine 1) if the composting process is complete or simply stalled out because of lack of nitrogen and 2) whether the compost, when applied to the soil, will act as a source of nitrogen for the crop or become a sink causing the crops to starve for nitrogen.

Your compost will tend to release available N for crop use. Your C:N ratio was 11

Ag Index

The Ag Index is the sum of nutrients N, P & K divided by the sum of non-nutrient salts Na & Cl. It provides an indication of whether your compost is a reasonable source of nutrients or primarily a source of organic matter for your soil. Your Ag index was q Your compost is a good source of nutreints and organic matter

Electrical Conductivity/Salinity

Electrical Conductivity is a convenient way to evaluate the soluble salts or salinity of a compost. High salinity is damaging to plants. Medium: best to dilute 1:2 to 1:5 for most applications The EC of your Compost was 4.8

Page 4